


IMAGES IN CARDIOLOGY

Platypnea-Orthodeoxia Syndrome

Insights of Mechanism From Imaging

Shohei Yoshida, MD,* Sawa Nambu, MD,† Takao Matsubara, MD,† Toshihiko Yasuda, MD,† Kenji Miwa, MD,† Masaru Inoue, MD,† Ryota Teramoto, MD,† Hirofumi Okada, MD,† Hounin Kanaya, MD,† Makoto Tsubota, MD,† Tetsuo Konno, MD,* Kenshi Hayashi, MD,* Masa-aki Kawashiri, MD,* Masakazu Yamagishi, MD*

Kanazawa, Japan



From the *Division of Cardiovascular Medicine, Kanazawa University Graduate School of Medicine, Kanazawa, Japan; †Division of Cardiology, Ishikawa Prefectural Central Hospital, Kanazawa, Japan; and the ‡Division of Cardiovascular Surgery, Ishikawa Prefectural Central Hospital, Kanazawa, Japan.
Manuscript received February 4, 2013; accepted February 19, 2013.

A 62-year-old man was admitted to our hospital because of cerebral infarction. Interestingly, oxygen saturation declined from 92% in supine to 78% in the upright position. Transesophageal echocardiography revealed the atrial septal defect (ASD) 10 mm in diameter. Although color Doppler demonstrated only left-to-right shunt, contrast imaging revealed mild bidirectional shunt in the supine position (**A and B**, [Online Videos 1 and 2](#)). Surprisingly, massive right-to-left shunt occurred in the upright position (**C and D**, [Online Videos 3 and 4](#)).

Multislice computed tomography in the upright position demonstrated the right-to-left shunt through the ASD (**E**, **arrowhead**) and the compressed right atrium by elongated aorta that might enhance right-to-left shunt (**E**, **arrows**). There was no evidence of pulmonary hypertension; thus, the patient was diagnosed as platypnea-orthodeoxia syndrome associated with ASD, which was confirmed on surgery to be closed (**F**, **arrow**). Ao = aorta; LA = left atrium; LV = left ventricle; RA = right atrium.